

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329279627>

Suicidal Thoughts, Attempts and Motives Among University Students in 12 Muslim-Majority Countries

Article in *Psychiatric Quarterly* · November 2018

DOI: 10.1007/s11126-018-9613-4

CITATIONS

0

READS

289

20 authors, including:



Mehmet Eskin

Koc University

86 PUBLICATIONS 1,512 CITATIONS

[SEE PROFILE](#)



Mohsen Rezaeian

Rafsanjan University of Medical Sciences

251 PUBLICATIONS 1,262 CITATIONS

[SEE PROFILE](#)



Ahmed M Abdel-Khalek

Alexandria University

330 PUBLICATIONS 4,911 CITATIONS

[SEE PROFILE](#)



Hacer Harlak

AydIn Adnan Menderes University

25 PUBLICATIONS 237 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Global Burden of Disease [View project](#)



Economic Burden of mental health in Pakistan [View project](#)

Suicidal Thoughts, Attempts and Motives Among University Students in 12 Muslim- Majority Countries

**Mehmet Eskin, Fadia AlBuhairan,
Mohsen Rezaeian, Ahmed M. Abdel-
Khalek, Hacer Harlak, Mayssah El-
Nayal, Nargis Asad, Aqeel Khan**

Psychiatric Quarterly

ISSN 0033-2720

Psychiatr Q

DOI 10.1007/s11126-018-9613-4

Volume 84 • Number 3 • September

**ONLINE
FIRST**

Psychiatric Quarterly

Available
online
www.springerlink.com

 Springer

11126 • ISSN 0033-2720
84(3) 271–406 (2013)

 Springer

Your article is protected by copyright and all rights are held exclusively by Springer Science+Business Media, LLC, part of Springer Nature. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Suicidal Thoughts, Attempts and Motives Among University Students in 12 Muslim-Majority Countries

Mehmet Eskin¹ · Fadia AlBuhairan^{2,3} · Mohsen Rezaeian⁴ · Ahmed M. Abdel-Khalek⁵ · Hacer Harlak⁶ · Mayssah El-Nayal⁷ · Nargis Asad⁸ · Aqeel Khan⁹ · Anwar Mechri¹⁰ · Isa Multazam Noor¹¹ · Motasem Hamdan¹² · Ulker Isayeva¹³ · Yousef Khader¹⁴ · Alaa Al Sayyari¹⁵ · Albaraa Khader¹⁴ · Bahareh Behzadi⁴ · Cennet Şafak Öztürk⁶ · Laifa Annisa Hendarmin¹⁶ · Murad Moosa Khan⁸ · Salam Khatib¹⁷

Published online: 29 November 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

There is a scarcity of research on suicidal phenomena in the Muslim world. Therefore, this study aimed at investigating the self-reported prevalence of suicidal thoughts, attempts and motives in 12 Muslim countries. A total of 8417 (54.4% women) university students were surveyed by means of a self-report questionnaire. Overall, 22% of the participants reported suicidal ideation and 8.6% reported attempting suicide. The odds of suicidal thoughts were elevated in Azerbaijan, Indonesia and Saudi Arabia, while reduced ORs were recorded in Egypt, Jordan, Lebanon and Malaysia. While odds of suicide attempts were high in Azerbaijan, Palestine and Saudi Arabia reduced odds ratios (OR) were detected in Indonesia, Iran, Jordan, Lebanon, Malaysia and Tunisia. Taking drugs and using a sharp instrument were the two most frequently used methods to attempt suicide. Only 32.7% of attempts required medical attention. Escape motives were endorsed more than social motives by participants who attempted suicide. Suicidal behaviors were more frequent in women than in men. Compared to men, fewer attempts by women required medical attention. Moreover, our results show that making suicide illegal does not reduce the frequency of suicidal behavior. Results from this comparative study show that suicidal thoughts and attempts are frequent events in young adults in countries where religious scripture explicitly prohibit suicide and the frequencies of nonfatal suicidal behavior show large variation in nations adhering to the same religion.

Keywords Suicide ideation · Suicide attempt · Suicidal motives · Young adults · Muslim world

Except the first author, the first twelve authors are by random order and the last seven authors are by alphabetical (first name) order.

✉ Mehmet Eskin
meskin@ku.edu.tr; meskin48@gmail.com

Extended author information available on the last page of the article

Introduction

Suicidal phenomena are a global public health concern and include suicidal thoughts, plans, attempts and deaths. Suicide is the cause of many premature deaths. It is a tragedy not only for the deceased but also for families, friends, neighbors and society. According to World Health Organization (WHO), 800,000 people die by suicide annually and it is the second leading cause of death among 15–29 year olds [1]. Seventy eight percent of all global suicides occur in low-middle income countries and the data indicate that there are 20 attempts for every suicidal death [1]. This means globally 16,000,000 individuals per annum make an attempt to kill themselves. Suicidal phenomena are therefore without a doubt significantly contributing to the burden on national health care systems [2]. For instance, the annual cost of suicide and suicide attempts are estimated to be 93.5 billion dollars in the United States [3].

Suicidal behavior is a multifaceted phenomenon that involves biological, genetic, psychological, social and cultural factors. As the WHO data suggest suicidal mortality peaks at 15–29 age group, corresponding roughly to the years during university education. Research indicates that factors associated with suicidal behavior in young people include mental health problems [4], alcohol and substance abuse [5], sexual minority status [6, 7], familial factors such as parental loss, discord and separation [8–10], academic stress [10, 11], economic difficulties [12] and low social support [13, 14].

All religions prohibit suicide with varying degrees of severity. Among the three Abrahamic religions, Islam has the most severe prohibitive attitude towards self-killing. In Judeo/Christian tradition, objections to suicide are based on the sanctity of life and the sixth commandment “Thou shalt not kill” [15]. In Islam, the objection against suicide comes as a word of God. The 4th surah of verse 29 of the Koran states “And do not kill yourselves [or one another]. Indeed, Allah is to you ever Merciful.” [16]. Hence, low official suicide mortality rates in Muslim majority nations are often attributed to the religion’s attitude. However, despite official statistics, the research suggests that the frequencies of nonfatal suicidal behaviors may indeed be high. In a number of cross-national comparisons, self-reported suicide attempts were significantly more frequent in Turkish adolescents and young adults than in their Austrian [17], Swedish [18] and Slovak [19] counterparts. Further, recently Eskin and colleagues [20] have shown that in 5572 university students from 12 countries, 29% reported suicidal ideation and 7% said that they attempted to kill themselves. Findings from this research showed that suicide attempts were more likely to be in the samples from Muslim majority countries such as Jordan, Palestine, Saudi Arabia and Turkey yet suicidal thoughts were less likely to be in those samples.

There is a dearth of scientific investigations on suicidal behavior in the Muslim majority countries and the existing ones are at the descriptive level [21]. The assumption that suicidal phenomena are rare in Muslim countries may be problematic for several reasons. First, official suicide rates are prone to be under and/or misreported because of the major stigma that resides with this issue [22]. Second, the Muslim majority world is not a homogenous entity. Muslim countries vary in their legal systems, social structures, values and attitudes, economic development, educational systems, indigenous cultures, ethnic composition, legal status of suicide and so on. As reported by the World Health Organization [1], 78% of all suicidal mortality occur in low-middle income countries, and the majority of Muslim nations lie within this category. From a methodological point of view, asking participants whether they have considered, and/or attempted suicide may sometimes provide a vague picture in what attempting suicide clearly entails. Indeed, in a recent paper Eskin, and colleagues [20] have called for an in-depth investigations of suicidal behavior especially in Muslim majority countries.

Bearing these considerations in mind, we designed this study to explore the aspects of suicidal thoughts, attempts, methods used and motives for suicide attempts in samples of university students from 12 Muslim majority countries. To the best of our knowledge, this is the first large scale comparative empirical study of suicidal behavior in the Islamic world. From a methodological perspective, cross-national comparative studies require that participants are comparable [23]. University students are similar in age and level of education, intellectual capacity, media exposure and stress levels. Therefore, we have chosen to study aspects of suicidal ideation and attempts in university students.

Methods

Participants

Participants in this cross-sectional study were 8417 university students from 12 Muslim majority countries (Azerbaijan, Egypt, Indonesia, Iran, Jordan, Lebanon, Malaysia, Pakistan, Palestine, Saudi Arabia, Tunisia and Turkey). The number of participants refusing to take part in the study were 10 in Malaysia and Turkey, 40 in Palestine, 115 in Iran, 145 in Tunisia, 20 in Egypt, and 28 in Pakistan. No one refused participation in Azerbaijan, Indonesia, Lebanon and Jordan yet it was not documented in Saudi Arabia. Seven questionnaires in Egypt and Jordan, 23 in Azerbaijan and Malaysia, 12 in Palestine, 14 in Iran, 48 in Tunisia, 40 in Turkey, 170 in Lebanon and 462 questionnaires in Saudi Arabia and no questionnaires in Indonesia and Pakistan were discarded due to incomplete information. Table 1 shows the sociodemographic characteristics of the participants. Overall, there were more women (54.4%) than men (45.6%) in our sample, $\chi^2 = 64.47$, $df = 1$, $p < 0.001$ (15 students did not report their gender). The gender distribution of students differed significantly by country, $\chi^2 = 104.86$, $df = 11$, $p < 0.001$; there were more women than men in the Indonesian, Palestinian, Saudi Arabian and Turkish samples, while the opposite was true for Malaysian sample.

Table 1 Demographic characteristics of participants according to country

Country	N	Gender				Age		Mother died		Father died		Par.Sep.		# of siblings	
		Women		Men		M	SD	n	%	n	%	n	%	M	SD
		n	%	n	%										
Azerbaijan	711	356	50.9	344	49.1	20.1	2.2	18	2.6	39	5.8	35	5.7	1.8	1.0
Egypt	653	328	50.3	324	49.7	20.3	1.2	10	1.5	64	9.9	27	4.7	2.6	1.4
Indonesia	300	208	69.3	92	30.7	19.8	1.0	9	3.0	9	3.0	10	3.4	2.2	1.3
Iran	700	350	50.0	350	50.0	22.1	4.0	13	1.9	41	5.9	10	1.5	2.6	1.8
Jordan	700	350	50.0	350	50.0	21.1	1.6	17	2.4	35	5.0	23	3.5	5.0	2.7
Lebanon	706	361	51.1	345	48.9	19.8	1.6	6	0.8	23	3.3	39	5.8	2.7	1.7
Malaysia	560	267	47.7	293	52.3	26.8	3.3	13	2.3	12	2.1	16	2.9	1.9	1.1
Pakistan	700	376	53.7	324	46.3	21.1	2.2	21	3.3	50	7.1	15	2.4	3.3	2.3
Palestine	793	474	60.0	316	40.0	20.4	3.1	14	1.8	48	6.1	16	2.2	5.3	2.1
Saudi Arabia	1137	711	62.5	426	37.5	22.0	3.6	17	1.5	88	7.7	96	9.4	5.8	2.9
Tunisia	707	355	50.2	352	49.8	21.0	2.0	9	1.3	30	4.2	27	4.1	2.6	1.4
Turkey	750	433	57.7	317	42.3	20.6	2.1	10	1.3	29	3.9	45	6.4	2.1	1.7
Total	8417	4569	54.4	3833	45.6	21.3	3.1	157	1.9	468	5.6	359	4.6	3.4	2.4

The mean age of the total sample was 21.28 (SD = 3.10) years with a range from 17 to 45 years. The mean ages differed significantly among countries, $F_{(11, 8405)} = 305.48, p < 0.001$. Indonesian and the Lebanese samples were the youngest and the Malaysian sample was the oldest. National samples differed significantly in the numbers of siblings, $F_{(11, 8405)} = 413.50, p < 0.001$. The Pakistani, Jordanian, Palestinian and the Saudi Arabian samples had the largest number of siblings, while the Azerbaijani, Malaysian, Turkish and the Indonesian samples had the smallest number of siblings.

While there were no differences between samples in terms of maternal death, $\chi^2 = 18.78, df = 11, p > 0.05$, they differed significantly from one another in paternal death, $\chi^2 = 66.90, df = 11, p < 0.001$. The largest percentages of Egyptian, Pakistani, Palestinian and Saudi Arabian but the smallest percentages in Indonesian, Lebanese, Malaysian and Turkish samples reported having gone through paternal death. Similarly, samples differed from one another in parental separation, $\chi^2 = 99.08, df = 11, p < 0.001$. Students from Saudi Arabia, Turkey, Lebanon and Azerbaijan reported highest numbers of parental separation and, those from Iran, Palestine, Pakistan, and Malaysia reported the lowest numbers.

We have retrieved age standardized suicide rates (per 100,000 population) for the countries included in the study from the World Health Organization's data base for the year 2015 [24]. The suicide rates were as follows for men, women and both, respectively: Azerbaijan (5.30, 1.10, 3.10), Egypt (4.50, 1.80, 3.10), Indonesia (4.50, 1.60, 3.00), Iran (4.20, 2.90, 3.60), Jordan (5.30, 2.40, 3.90), Lebanon (4.00, 2.00, 3.10), Malaysia (9.50, 3.40, 6.50), Pakistan (2.50, 2.40, 2.50), Palestine (NA), Saudi Arabia (5.50, 2.20, 3.90), Tunisia (6.70, 4.10, 5.40) and Turkey (12.60, 4.70, 8.60).

Instrument

A self-administered questionnaire was used to collect the data which contained questions and measures about aspects of nonfatal suicidal behavior, religious affiliation and strength of religious belief, knowledge about affiliated religion's attitude to suicide, negative life-events, attitudes towards suicide, social support, self-construal, religious coping, motives for suicide attempts and impacts of suicide attempts on personal-social relationships. In this article, the prevalence of suicidal behavior, and methods and motives for suicide attempts are reported. (In order to keep a clear focus, the associations of suicidal behavior to religion, suicidal attitudes, negative life-events and social support, and self-construal will be reported in separate papers). The questionnaire was translated into home languages (Arabic, Azerbaijani, Indonesian, Malay, Persian, Urdu and Turkish) by using the parallel blind method [25].

Suicidal Behavior

Two yes-no questions (Have you ever thought of killing yourself? and Have you ever attempted to kill yourself?) were administered on life-time suicide ideation and attempts.

Self-Rated Frequency and Severity of Suicidal Behavior

The frequency and severity of suicidal thoughts and attempts were asked on 5-point scales ranging from 1 (only once/not at all serious) to 5 (more than five times/very serious).

Time of and Desire to Die in Suicide Attempts

The time of and the intensity of desire to die in the latest attempts were asked. The responses were obtained on a 5-point scale ranging from 1 (1 year ago) to 5 (five or more years ago) for the former question and on a 5-point scale ranging from 1 (none) to 5 (very strong) for the latter.

Method of Suicide Attempts

The methods used to attempt suicide were enquired with a list of 10 methods. Participants could choose more than one method if they had more than one suicide attempt. Since fewer persons attempted suicide by burning ($n = 7$) and natural gas ($n = 7$) they were collapsed into the other methods category (see Table 4).

Medical Attention

To determine whether medical attention was required for an attempt or not, participants who attempted suicide were asked: What happened after the latest attempt? They responded to this question by choosing one of the four alternatives: (1) Nothing happened because it was not serious. (2) I was conscious, and they took me to the hospital and discharged me after a few hours. (3) I was conscious, and they took me to the hospital and I stayed in the hospital for a day or more for treatment. (4) I was unconscious, and they took me to the hospital and I stayed in the hospital for a day or more for treatment.

Attempts in the first alternative were denoted as “attempts requiring no medical attention” and those in the second to fourth categories as “attempts requiring medical attention”.

Motives for Suicide Attempts

Nine possible reasons for attempting suicide [26] were presented to participants and asked to indicate how important was each of them for their latest attempt on 5-point Likert scales ranging from “not at all important = 1” to “extremely important = 5”. To group the reasons, a principal component analysis with varimax rotation was employed. Results of the analysis revealed two factors which explained 61.26% of the total variance. Six items loaded on the first factor (1. To find out whether someone really loved me or not; 2. To try to influence someone or get them to change their mind; 3. To show how much I loved someone; 4. To make people sorry for the way they have treated me; 5. To frighten or get someone back; 6. To seek help from someone and to make people understand how desperate I was feeling). The factor loadings ranged from 0.62 to 0.82 and this factor was named as social motives ($\lambda = 3.75$; attributable variance = 41.65). Three items loaded on the second factor (1. To get relief from a terrible state of mind; 2. To escape for a while from an impossible situation; 3. To die). Factor loadings ranged from 0.64 to 0.85 and this factor was named as escape motives ($\lambda = 1.77$; attributable variance = 19.62). The internal consistency coefficient (Cronbach's α) for social motives factor was 0.86 and it was 0.65 for escape motives factor. Two factor scores were computed by summing up individual responses on the items loading on the respective factor and dividing this sum by the number of items corresponding to that factor. Hence, the possible range of factor scores was from 1 to 5. Higher scores indicate greater amount of factor content.

Demographics

Students were asked about their gender, age, number of siblings, parental loss and separation.

Procedure

The questionnaire and the study protocol were first prepared by the principal investigator (M. Eskin) in English. The principal investigator tracked the researchers through their publication records and invited them for collaboration via e-mail. The mail explained the background and the method (sample and the questionnaire) of the study. The researchers are psychologists, psychiatrists, epidemiologists, psychological counsellors and public health specialists. Four of the researchers (A. Mechri, H. Harlak, M. Hamdan, Y. Khader) collaborated in a previous international research project on suicide with the principle investigator.

Ethical or IRB approval was obtained in all study sites. Except for Saudi Arabia where data collection was done via a web-based survey, a paper and pencil questionnaire were used in the remaining study sites. On the first page of the questionnaire it was highlighted that the study was anonymous. The students were reminded that they did not need to provide personal details. The name, telephone number and e-mail address of the local investigator were provided on the first page of the questionnaire for participants who might have had personal concerns over the questions.

No adverse effects were observed during data collection. Site (country) researchers were sent a follow up researcher survey about the practicalities of data collection and the legal status of suicide. According to the responses to the researcher survey, the site researchers reported having no difficulty in obtaining IRB approval and permission from the administration of universities for collecting the data. Attempting suicide was illegal in Jordan, Malaysia, Pakistan, Palestine and Saudi Arabia according to the responses to the survey.

Statistical Analyses

Data were analyzed by using the SPSS-19 for windows. Percentages of suicidal thoughts and attempts, suicide methods and demographic characteristics of participants were calculated by country and gender. First, twelve countries and nine suicide methods were coded as dummy variables. The associations of country, gender, age, parental separation-loss and number of siblings were tested by calculating the odds ratios (OR) for suicidal thoughts and attempts by using logistic regression and Chi-square procedures. In calculating country ORs for suicidal behavior, age, gender and number of siblings were taken as covariates. The ORs between two dichotomous variables were calculated by Chi-square procedure. Group means were compared by means of *t*-tests and one-way analysis of variance (one-way ANOVA). Associations between two continuous variables were tested by Pearson product moment correlation coefficient and the associations between continuous and categorical variables were tested by point-biserial correlation coefficient procedures. Countries with legal sanctions against suicide and those without legal sanction were dichotomized into two groups. The difference between the two groups in country suicide rates ($n = 11$) was tested by Mann-Whitney U test procedure. Odds ratios between the legal status of suicide dichotomy and suicidal thoughts, attempts and medical attention were calculated by Chi-square procedure. The differences between the two groups in self-rated frequency and seriousness of suicidal thoughts and attempts, and timing of and intensity of death desire expressed in suicide attempts were tested by *t*-test procedure.

Results

Distribution and Characteristics of Suicidal Behavior

Table 2 displays the numbers, percentages, and mean frequency, severity, time and intensity of death desire in suicidal behavior by country. Overall, 1825 participants (22.1%) reported having had thoughts of killing themselves and 712 students (8.6%) reported having made at least one attempt to kill themselves.

Table 3 presents the country and gender odds ratios for suicide ideation and attempts. The odds for suicidal thoughts were significantly larger for participants from Azerbaijan, Indonesia and Saudi Arabia, while for the participants from Egypt, Jordan, Lebanon and Malaysia they were significantly smaller. The odds for suicidal attempts were significantly larger for

Table 2 Prevalence, frequency, severity of suicidal ideation and attempts

Country	Suicidal ideation						Suicidal attempt									
	N	%	Frequency		Severity		N	%	Frequency		Time of attempt		Desire to die			
			M	SD	M	SD			M	SD	M	SD	M	SD		
Azerbaijan	195	31.1	2.4	1.6	2.9	1.2	129	20.5	2.2	1.5	3.2	1.3	2.7	1.6	3.0	1.2
Egypt	114	17.5	2.5	1.5	2.4	1.3	46	7.1	2.2	1.6	2.9	1.3	2.8	1.6	3.1	1.4
Indonesia	95	31.7	2.0	1.5	1.3	0.8	12	4.0	1.3	1.2	1.8	1.0	2.7	1.9	2.5	0.9
Iran	160	22.9	3.1	1.6	2.1	1.3	36	5.1	1.8	1.1	3.1	1.3	2.5	1.6	3.1	1.3
Jordan	87	12.4	2.6	1.5	2.5	1.3	39	5.6	2.1	1.1	3.0	1.3	2.7	1.5	2.9	1.3
Lebanon	87	12.3	2.4	1.5	2.1	1.3	36	5.1	1.8	1.2	3.0	1.5	2.9	1.6	2.7	1.1
Malaysia	5	0.9	1.8	0.4	2.2	0.4	5	0.9	2.0	0.7	2.6	0.5	2.8	0.4	3.4	0.9
Pakistan	152	21.7	2.9	1.7	2.9	1.4	48	6.9	1.9	1.1	3.3	1.4	2.8	1.6	3.3	1.1
Palestine	174	23.6	2.7	1.6	2.2	1.3	125	17.6	2.1	1.3	2.7	1.4	2.2	1.3	2.6	1.3
Saudi Arabia	439	38.7	3.2	1.6	2.3	1.3	152	13.4	2.0	1.4	2.9	1.3	2.9	1.7	3.5	1.2
Tunisia	136	19.2	2.7	1.6	2.3	1.2	35	5.0	1.6	1.0	2.9	1.5	3.6	1.8	2.9	1.3
Turkey	181	24.1	2.8	1.7	2.8	1.2	49	6.5	2.2	1.5	3.6	1.1	2.6	1.5	3.3	1.1
Total	1825	22.1	2.8	1.6	2.4	1.3	712	8.6	2.0	1.3	3.0	1.3	2.8	1.6	3.1	1.2
χ^2 or F	465.17*		7.43*		14.63*		319.99*		1.30		2.90*		1.74		3.51*	

* All chi-square and F values are significant $p < 0.01$

Orderings of country frequency, seriousness, timing and death desire ratings from the lowest to the highest mean values:

Suicide Ideation:

Frequency: Malaysia, Indonesia, Lebanon = Azerbaijan, Egypt, Jordan, Palestine = Tunisia, Turkey, Pakistan, Iran, Saudi Arabia

Seriousness: Indonesia, Iran = Lebanon, Malaysia = Palestine, Saudi Arabia = Tunisia, Egypt, Jordan, Turkey, Azerbaijan = Pakistan

Suicide Attempts:

Frequency: Indonesia, Tunisia, Iran = Lebanon, Malaysia = Pakistan, Saudi Arabia, Jordan = Palestine, Azerbaijan = Egypt = Turkey

Severity: Indonesia, Malaysia, Palestine, Egypt = Saudi Arabia = Tunisia, Jordan = Lebanon, Iran, Azerbaijan, Pakistan, Turkey

Time of: Palestine, Iran, Turkey, Azerbaijan = Indonesia = Jordan, Egypt = Malaysia = Pakistan, Lebanon = Saudi Arabia, Tunisia

Desire to die: Indonesia, Palestine, Lebanon, Jordan = Tunisia, Azerbaijan, Egypt = Iran, Pakistan = Turkey, Malaysia, Saudi Arabia

Table 3 Country and gender odds ratios for suicidal ideation and attempts

Countries	Suicidal thoughts				Suicide attempts			
	Country		Gender (women = 1; men = 0)		Country		Gender (women = 1; men = 0)	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Azerbaijan	1.72*	(1.43 — 2.07)	2.05*	1.40 — 2.98	3.61*	(2.88 — 4.52)	2.16*	1.39 — 3.36
Egypt	0.73*	(0.59 — 0.90)	3.18*	1.96 — 5.16	0.80	(0.58 — 1.09)	2.71*	1.33 — 5.54
Indonesia	1.52	(1.18 — 1.96)	0.81	0.47 — 1.38	0.39*	(0.22 — 0.70)	2.01	0.41 — 9.83
Iran	1.19	(0.99 — 1.44)	0.96	0.66 — 1.39	0.64**	(0.45 — 0.90)	0.38**	0.18 — 0.81
Jordan	0.43*	(0.34 — 0.55)	1.08	0.65 — 1.78	0.53*	(0.38 — 0.75)	0.93	0.46 — 1.88
Lebanon	0.44*	(0.35 — 0.55)	2.60*	1.55 — 4.38	0.51*	(0.36 — 0.72)	1.71	0.83 — 3.51
Malaysia	0.03*	(0.01 — 0.08)	2.37	0.21 — 27.27	0.14*	(0.06 — 0.35)	2.39	0.21 — 27.43
Pakistan	0.98	(0.81 — 1.19)	1.43	0.97 — 2.12	0.77	(0.57 — 1.05)	1.17	0.61 — 2.23
Palestine	0.90	(0.75 — 1.09)	1.10	0.75 — 1.62	2.12*	(1.69 — 2.65)	1.16	0.74 — 1.83
Saudi Arabia	2.90*	(2.50 — 3.37)	1.26	0.96 — 1.64	1.69*	(1.37 — 2.09)	2.03*	1.31 — 3.15
Tunisia	0.87	(0.71 — 1.06)	1.69**	1.14 — 2.53	0.57*	(0.40 — 0.80)	2.79*	1.29 — 6.03
Turkey	1.15	(0.96 — 1.38)	1.46**	1.01 — 2.10	0.74	(0.55 — 1.00)	1.14	0.60 — 2.17

df= 1 for all; * Wald statistics are significant $p < 0.01$; ** Wald statistics are significant $p < 0.0.5$

participants from Azerbaijan, Palestine and Saudi Arabia yet were significantly smaller for participants from Indonesia, Iran, Jordan, Lebanon, Malaysia and Tunisia.

Table 2 shows that while Pakistani, Iranian and Saudi Arabian students reported having had the most frequent thoughts of suicide, Malaysian, Indonesian, Lebanese and Azerbaijani students reported the least frequent suicidal thoughts. Turkish, Azerbaijani and Pakistani students rated their suicidal thoughts as the most serious yet Indonesian, Iranian and Lebanese students as the least serious. The most frequent suicide attempters were from Azerbaijani, Egyptian and Turkish samples, least frequent attempters were from Indonesian, Tunisian, Iranian and Lebanese samples. While Azerbaijani, Pakistani and Turkish suicide attempters rated their attempts as the most serious, Indonesian, Malaysian and Palestinian attempters the least serious. Palestinian, Iranian, and Turkish students reported the most recent attempts yet Lebanese, Saudi Arabian and Tunisian students reported the most earlier attempts. Pakistani, Turkish, Malaysian and Saudi Arabian attempters expressed the intense desire to die at most, Indonesian, Palestinian and Lebanese attempters expressed the same motive at least.

In the total sample, suicidal thoughts were significantly more common in women ($B = 0.42$, $Wald_{(1)} = 52.47$, $p < 0.001$, $OR = 1.52$, $95\%CI = 1.36 — 1.70$) than men. Table 3 shows also that the odds for suicidal thoughts were significantly larger for women than men in Azerbaijan, Egypt, Lebanon, Tunisia and Turkey. In the whole sample, significantly more women than men reported attempting suicide ($B = 0.44$, $Wald_{(1)} = 24.17$, $p < 0.001$, $OR = 1.54$, $95\%CI = 1.30 — 1.87$). Table 3 displays that the odds for suicide attempts were significantly larger for women than men in Azerbaijan, Egypt, Saudi Arabia and Tunisia yet it was significantly smaller in Iran.

Significantly more participants who were younger ($B = -0.08$, $Wald_{(1)} = 56.77$, $p < 0.001$, $OR = 0.93$, $95\%CI = 0.91 — 0.95$), whose mothers died ($\chi^2 = 5.84$, $p < 0.05$, $OR = 1.54$, $95\%CI = 1.08 — 2.18$), whose parents separated ($\chi^2 = 53.08$, $p < 0.001$, $OR = 2.25$, $95\%CI = 1.80 — 2.81$), and those with larger number of siblings ($B = 0.04$, $Wald_{(1)} = 12.94$, $p < 0.001$, $OR = 1.04$, $95\%CI = 1.02 — 1.06$) reported suicide ideation.

Similarly, significantly more participants who were younger ($B = -0.10$, $Wald_{(1)} = 34.68$, $p < 0.001$, $OR = 0.91$, $95\%CI = 0.88 - 0.94$), whose mothers ($\chi^2 = 13.90$, $p < 0.001$, $OR = 2.22$, $95\%CI = 1.44 - 3.40$) and fathers ($\chi^2 = 8.18$, $p < 0.01$, $OR = 1.52$, $95\%CI = 1.14 - 2.04$) died, whose parents separated ($\chi^2 = 39.19$, $p < 0.001$, $OR = 2.45$, $95\%CI = 1.83 - 3.27$) and those with larger number of siblings ($B = 0.06$, $Wald_{(1)} = 16.93$, $p < 0.001$, $OR = 1.08$, $95\%CI = 1.06 - 1.09$) stated attempting suicide.

Methods Used to Attempt Suicide

Table 4 displays the number and percentages of suicide attempts by country. The two most frequently used methods for attempting suicide were by taking chemicals, i.e. pills, and using a sharp instrument, e.g. knife. As the table shows, methods used for attempting suicide differ significantly between countries, $\chi^2_{(88)} = 248.54$, $p < 0.001$.

Methods used for attempting suicide differed significantly between genders, $\chi^2_{(8)} = 73.97$, $p < 0.001$. Methods of hanging ($\chi^2_{(1)} = 15.22$, $p < 0.001$, $OR = 0.33$, $95\%CI = 0.19 - 0.59$), using fire arms ($\chi^2_{(1)} = 14.62$, $p < 0.001$, $OR = 0.17$, $95\%CI = 0.06 - 0.47$), jumping ($\chi^2_{(1)} = 14.15$, $p < 0.001$, $OR = 0.35$, $95\%CI = 0.20 - 0.62$), and drowning ($\chi^2_{(1)} = 3.92$, $p < 0.05$, $OR = 0.42$, $95\%CI = 0.18 - 1.02$) were significantly more likely to be used by men but taking chemicals such as pills ($\chi^2_{(1)} = 37.21$, $p < 0.001$, $OR = 3.13$, $95\%CI = 2.15 - 4.56$) and using a sharp instrument ($\chi^2_{(1)} = 4.48$, $p < 0.05$, $OR = 1.58$, $95\%CI = 1.03 - 2.40$) were more likely to be used by women.

Methods of suicide attempts differed significantly according to whether an attempt required medical attention or not, ($\chi^2_{(1)} = 21.04$, $p < 0.01$). Significantly more attempts involving taking chemicals such as pills required medical attention ($\chi^2_{(1)} = 12.77$, $p < 0.001$, $OR = 1.88$, $95\%CI = 1.33 - 2.66$) but significantly fewer attempts involving a sharp instrument ($\chi^2_{(1)} = 7.90$, $p < 0.01$, $OR = 0.53$, $95\%CI = 0.33 - 0.83$) and other methods ($\chi^2_{(1)} = 4.94$, $p < 0.05$, $OR = 0.48$, $95\%CI = 0.25 - 0.93$) required medical attention.

Medical Attention

Of 712 suicide attempters, 612 (85.96%) responded to the question about what happened after the latest attempt. An overwhelming majority ($n = 412$, 67.30%) said "nothing happened because it was not serious". One hundred and eleven (18.10%) said they were conscious and taken to hospital and discharged after a few hours. Fifty-two (8.50%) said they were conscious and taken to hospital and stayed there for a day or more for treatment. Finally, 37 (6.00%) said they were unconscious and taken to hospital and stayed there for a day or more for treatment after the attempt. Thus, suicide attempts by 200 (32.70%) participants required medical attention.

Comparison of participants whose attempts required medical attention to those whose attempts did not require medication attention revealed that fewer attempts by women (women $n = 108$, 27.8% versus men $n = 91$, 40.8%; $\chi^2 = 10.85$, $OR = 0.56$, $95\% CI = 0.40 - 0.79$) required medical attention however the two groups did not differ in age, number of siblings, parental separation and loss. Compared to participants whose attempts did not require medication attention, those whose attempts required medical attention rated their attempts as more serious ($M = 2.76$, $SD = 1.25$ versus $M = 3.57$, $SD = 1.28$, $t_{(607)} = 7.42$, $p < 0.001$, Cohen's $d = 0.64$) and gave higher death desire ratings ($M = 2.91$, $SD = 1.22$ versus $M = 3.57$, $SD = 1.16$, $t_{(603)} = 6.34$, $p < 0.001$, Cohen's $d = 0.55$). The two groups were similar in the frequency ratings and timing of their attempts.

Table 4 Methods used for attempting suicide by country

Countries	Hanging		Fire arms		Throwing		Jumping		Sharp Instr.		Drowning		Chemicals		Poisoning		Other	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Azerbaijan	10	8.3	3	2.5	4	3.3	12	10.0	21	17.5	9	7.5	41	34.2	6	5.0	14	11.7
Egypt	1	2.2	0	0.0	2	4.4	3	6.7	6	13.3	1	2.2	19	42.2	4	8.9	9	20.0
Indonesia	0	0.0	0	0.0	0	0.0	1	8.3	5	41.7	1	8.3	2	16.7	2	16.7	1	8.3
Iran	5	13.9	0	0.0	0	0.0	2	5.6	7	19.4	0	0.0	15	41.7	1	2.8	6	9.7
Jordan	5	13.5	0	0.0	0	0.0	5	13.5	6	16.2	3	8.1	12	32.4	1	2.7	5	13.5
Lebanon	4	11.1	4	11.1	3	8.3	3	8.3	5	13.9	0	0.0	15	41.7	1	2.8	1	2.8
Malaysia	0	0.0	3	60.0	2	40.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pakistan	2	4.2	1	2.1	2	4.2	3	6.3	16	33.3	2	4.2	13	27.1	7	14.6	2	4.2
Palestine	13	28.9	2	4.4	5	11.1	4	8.9	9	20.0	1	2.2	7	15.6	2	4.4	2	4.4
Saudi Arabia	10	6.6	2	1.3	3	2.0	13	8.6	46	30.3	4	2.6	56	36.8	2	1.3	16	10.5
Tunisia	2	5.7	1	2.9	0	0.0	5	14.3	6	17.1	1	2.9	18	51.4	0	0.0	2	5.7
Turkey	2	4.1	5	10.2	2	4.1	4	8.2	2	4.1	0	0.0	29	59.2	1	2.0	4	8.2
Total	54	8.7	21	3.4	23	3.7	55	8.9	129	20.8	22	3.5	227	36.6	27	4.4	62	10.0

While the odds for attempts requiring medical attention were significantly greater for the attempts being from Lebanon ($\chi^2(1) = 7.02, p < 0.01, OR = 2.44, 95\% CI = 1.24 - 4.80$), and Malaysia ($\chi^2(1) = 10.39, p < 0.01, OR = 0.1.03, 95\% CI = 1.01 - 1.05$), yet smaller for the attempts being from Saudi Arabia ($\chi^2(1) = 9.77, p < 0.01, OR = 0.51, 95\% CI = 0.33 - 0.78$).

Motives for Attempting Suicide

Participants who attempted suicide gave the highest ratings to the two escape related items (to escape for a while from an impossible situation; to get relief from a terrible state of mind) and to one item related to social motives (to make people sorry for the way they've treated me; frighten or get someone back). Table 5 presents the means and standard deviations for the two motives for attempting suicide factor scores by country. The countries significantly differed in the mean scores of escape and of social motives for attempting suicide. The highest mean of escape motives was noted in Saudi Arabian, Turkish and Iranian attempters yet the lowest were seen in Malaysian, Indonesian and Palestinian attempters. The highest mean of social motives was observed in Azerbaijani, Iranian and Pakistani attempters while the lowest were noted in Turkish, Saudi Arabian and Indonesian attempters.

The mean of escape motive scores for men ($M = 3.04, SD = 1.14$) and women ($M = 3.18, SD = 1.11$) were similar, $t(603) = 1.53, p > 0.05$ while the mean of social motive scores for men ($M = 2.61, SD = 1.11$) were significantly greater than the mean of social motive scores for women ($M = 2.30, SD = 1.17$), $t(602) = 3.25, p < 0.01$.

Table 5 Mean escape and social motives for attempting suicide by country

Countries	Escape motives		Social motives	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Azerbaijan	2.77	1.21	2.58	1.18
Egypt	3.03	1.18	2.42	1.15
Indonesia	2.56	0.92	2.24	1.37
Iran	3.83	1.00	2.78	1.35
Jordan	2.81	0.92	2.39	0.98
Lebanon	2.65	0.99	2.57	0.98
Malaysia	2.40	0.55	2.53	0.68
Pakistan	3.38	1.04	2.98	1.27
Palestine	2.59	0.95	2.48	1.14
Saudi Arabia	3.50	1.04	2.10	1.07
Tunisia	2.77	1.00	2.25	1.15
Turkey	3.56	0.94	2.08	1.01
Total	3.13	1.12	2.41	1.15
<i>F</i> values*	8.19		3.28	

* *df* for *F* values for escape motives = (11, 594) and for social motives = (11, 593). Both *F* values are significant at $p < 0.001$

Orderings of country escape and social motive factor scores from the lowest to the highest mean values:

Escape motives:

Malaysia, Indonesia, Palestine, Lebanon, Tunisia, Azerbaijan, Jordan, Egypt, Pakistan, Saudi Arabia, Turkey, Iran

Social motives:

Turkey, Saudi Arabia, Indonesia, Tunisia, Jordan, Egypt, Palestine, Malaysia, Lebanon, Azerbaijan, Iran, Pakistan

While self-hanging to attempt suicide was inversely associated with escape motives, $r = 0.10$, $p < 0.05$, taking chemicals (pills) was positively related to escape motives, $r = 0.12$, $p < 0.01$. Only self-poisoning was related to endorsing more social motives, $r = 0.09$, $p < 0.05$.

Iranian ($r = 0.16$, $p < 0.001$), Saudi Arabian ($r = 0.19$, $p < 0.001$) and Turkish ($r = 0.12$, $p < 0.01$) attempters endorsed escape motives to attempt suicide to a greater extent than others, yet Lebanese ($r = -0.10$, $p < 0.05$), Palestinian ($r = -0.13$, $p < 0.01$) and Azerbaijani ($r = -0.16$, $p < 0.001$) attempters obtained lower escape motive scores than others. Iranian ($r = 0.08$, $p < 0.05$) and Pakistani ($r = 0.15$, $p < 0.001$) attempters obtained higher social motive scores while Saudi Arabian ($r = -0.15$, $p < 0.001$) and Turkish ($r = 0.09$, $p < 0.05$) attempters had lower social motive scores than others.

There were statistically significant correlation coefficients between escape motives and, self-rated severity of attempts ($r = 0.21$, $p < 0.001$) and intensity of desire to die ($r = -0.31$, $p < 0.001$). The social motives were related only to the time of latest attempt ($r = -0.11$, $p < 0.01$).

Legal Status of Suicide

Age standardized suicide rates (per 100,000 population) for countries where suicide is illegal were not different than the rates for countries where there is no legal sanction against suicide (rates for both sexes $Z = -0.38$, $p > 0.05$; men $Z = -0.29$, $p > 0.05$; women $Z = 0.57$, $p > 0.05$). Suicidal thoughts were equally frequent in the countries where suicide is illegal and in those where there is no legal sanctions against suicide, ($\chi^2(1) = 0.39$, $p > 0.05$, OR = 1.03, 95% CI = 0.93 — 1.15) but suicide attempts were more frequent in the countries where suicide is illegal than in others, ($\chi^2(1) = 10.12$, $p < 0.01$, OR = 1.28, 95% CI = 1.10 — 1.50). Equal numbers of attempts in these two group of countries required medical attention, ($\chi^2(1) = 2.31$, $p > 0.05$, OR = 0.77, 95% CI = 0.55 — 1.08).

In countries where suicide is illegal participants reported having had suicidal thoughts more frequently ($M = 2.99$, $SD = 1.64$ versus 2.61, $SD = 1.62$, $t(1752) = 4.82$, $p < 0.001$) and expressed greater death desire in their attempts ($M = 3.23$, $SD = 1.26$ versus 3.00, $SD = 1.23$, $t(625) = 2.18$, $p < 0.05$) than their counterparts in countries where there is no legal sanction against suicide. The two groups were similar in seriousness of both suicidal thoughts and attempts and, frequency and timing of the attempts.

Discussion

This study revealed some aspects of nonfatal suicidal behavior in university students from 12 Muslim majority countries. Our findings, in line with some previous work, suggest that, suicidal thoughts (22.1%) and attempts (8.6%) are frequent events in young adults in Muslim majority countries despite explicit condemnation of suicide by Islam [17, 19, 27, 28]. These percentages speak for themselves in that so many gifted young women and men have considered and attempted to kill themselves despite a potential for a happy and productive life ahead. Mental disorders are associated with suicidal mortality [29, 30] and the burden of disease estimates indicate that the burden of mental disorders is rising and highest in the 25–49 age group in the Eastern Mediterranean Region [31] from where the majority of samples in this study originated.

It is interesting to note that the rates of suicidal ideation and attempts are comparable to or higher than the rates observed in university students in The United States [32, 33] and China

[34]. Considering the attitude of Islam to suicide one would have expected to see lower rates of suicidal ideation and attempts in the Islamic world. In the last decade Muslim societies have faced serious challenges. Political instability, lack of freedom and liberties, unemployment, gender inequality, large youth population competing for limited resources and dissatisfaction with the existing regimes are the major challenges [35, 36]. The high suicidal thought and attempt rates might reflect these instable and threatening social conditions.

The results from this multinational study show clearly that the frequencies of suicidal behaviors vary across nations included in the study. Compared to the rest, significantly more students from Azerbaijan, Indonesia and Saudi Arabia reported having thought to kill themselves in contrast to significantly fewer students from Egypt, Jordan, Lebanon and Malaysia reported suicide ideation. Significantly more students from Azerbaijan, Palestine and Saudi Arabia reported attempting suicide in comparison to fewer students from Indonesia, Iran, Jordan, Lebanon and Malaysia. In a previous multinational study, Eskin and colleagues found that significantly more Palestinian and Saudi Arabian students said that they attempted to kill themselves [20]. In line with this finding, Palestine had the largest burden of mental disorders among the EMR countries [31].

In surveys, what does an affirmative response to a question on suicide attempt entail is not clear. To gauge the nature of suicidal thoughts and attempts in detail participants were asked about their perceptions of frequency, seriousness, timing and death desire in their attempts. Frequency and seriousness ratings of participants with suicidal thoughts from national samples differed significantly in their responses from one another. Saudi Arabian, Iranian, Pakistani and Turkish students reported having entertained such thoughts more frequently than others. Students having more frequent suicidal ideation rated their thoughts more serious ($r = 0.22$, $p < 0.001$). Pakistani, Azerbaijani, Turkish and Jordanian students reported their suicidal thoughts to be more serious than others. Turkish, Pakistani, Azerbaijani, Iranian students rated their attempts more serious than others. Saudi Arabian, Malaysian, Turkish, Pakistani students reported having had more death desire in their attempts than others. Participants with more frequent attempts rated their attempts as more serious ($r = 0.19$, $p < 0.001$), as more recent ($r = -0.26$, $p < 0.001$) and involving more death desire ($r = 0.14$, $p < 0.01$). Students with more serious attempts reported more death desire in their attempts ($r = 0.58$, $p < 0.001$).

A self-reported suicidal behavior may have multiple meanings. One way of gauging the nature of suicide attempts is to see whether an attempt required medical attention or not. According to our results, 67.3% of attempters said that nothing happened after the attempt because it was not serious. Thus, only 32.7% of attempts required some form of medical attention. Significantly more attempts in Malaysian and Lebanese in contrast to fewer attempts from Saudi Arabian samples required medical attention. Further, our results revealed that attempts requiring medical attention were rated as more serious and involving more death desire than the attempts not requiring medical attention. The reason for most attempts not requiring medical attention may be due to three reasons. First, the lack of or scantiness of mental health services might lie behind this observation. Another reason might be the mental stigma. It is noted that mental stigma is widespread among Muslims [37]. A third reason might be an overlap between non-suicidal self-injurious acts and suicide attempts. Non-suicidal self-injurious acts are far more frequent than suicidal attempts and constitute a major risk factor for suicide [38]. It is possible that participants with suicide attempts requiring medical attention perceived their attempts as more serious and expressed greater amounts of death desire than those whose attempts did not require medical attention.

Suicide methods vary across cultural and/or national [39] and racial groups [40]. The most frequent methods used for attempting suicide, in this study, were taking chemicals e.g. pills,

using a sharp instrument, jumping from a high place and hanging, and they showed a large variation across nations under scrutiny. The variation of methods by country reflects probably the ease at reaching these means. It is interesting to note that significantly more attempts involving taking chemicals required medical attention, but fewer attempts involving use of sharp instrument required medical attention. Self-injurious behaviors such as cutting and carving the skin [41] has a potential to evolve into suicidal attempt [42]. However, it is interesting to note that significantly more attempts involving taking chemicals required medical attention, but fewer attempts made use of sharp instrument required medical attention.

Suicidal behavior and method choice show a gendered pattern. Women report suicidal thoughts and attempts more often than men yet men kill themselves more often than women [43]. Further compared to men, women choose less lethal methods for suicide [44, 45]. In line with this point, the present study revealed that significantly more women than men reported suicidal ideation (in the total sample, Azerbaijani, Egyptian, Lebanese, Tunisian and Turkish samples) and attempts (in the total sample, Azerbaijani, Egyptian, Saudi Arabian and Tunisian samples) but more men than women reported attempting suicide in the Iranian sample. More men than women reported using more lethal methods (hanging, fire arms, jumping and drowning) however more women than men used less lethal methods (taking pills and using sharp instrument). Accordingly, fewer attempts by women than men required medical attention.

Peoples' motives for attempting suicide may vary across cultural, gender, age groups and so on. Consistent with previous work [26] suicide attempters in our study endorsed escape motives to a greater extent than social motives. Attempters with higher mean of escape motives rated their attempts more serious and reported higher levels of death desire in their attempts. While Iranian, Turkish, Saudi Arabian and Pakistani attempters had the highest mean of escape motives Indonesian, Palestinian, Lebanese and Tunisian attempters had the lowest mean of escape motives. While Pakistani, Iranian, Azerbaijani and Lebanese attempters had the highest mean social motives Turkish, Saudi Arabian, Indonesian and Tunisian attempters had the lowest mean of social motive scores.

The goal of any legal sanction against suicidal behavior, whether it is based on religious or secular grounds, should be to lower the frequencies of suicidal behavior thereby save lives. But does it function as intended? Our results show that prohibiting suicidal behavior by law does not work in an intended direction. Official suicidal mortality rates and the frequencies of self-reported suicidal thoughts in the countries with and without legal sanction against suicide were similar. On the contrary, self-reported suicide attempts were more frequent in the countries with legal sanction against suicide than in those with no such sanction. Moreover, participants in the countries where suicide is illegal reported having had more frequent suicidal thoughts and expressed greater death desire in their suicide attempts than their counterparts in countries where suicide is not forbidden by law. Making suicide illegal may aggravate the further stigmatization of an already stigmatized phenomenon. This in turn may function as an obstacle against help seeking and a hinder for social connectedness which are both shown to be associated with lower propensity for suicidal behavior [2, 46, 47].

Limitations

Although current findings shed some light on the variation and the nature of suicidal phenomena in university students in Muslim majority countries, they should be approached with caution for several reasons. First, we have no claim that our samples are representative of

the general populations of countries under scrutiny. The university students are representative of neither the young segments of countries nor the whole populations. Second, though anonymous, self-administered paper and pencil questionnaires may not be suitable for collecting data on culturally sensitive topics such as suicide. Web-based surveys may be more suitable for this purpose. Since data collection was done via a web-based survey, greater reports of suicidal ideation and attempts in Saudi Arabian sample and lower frequencies in Jordanian, Malaysian and Pakistani samples may be reflecting the effects of the medium of data collection. Third, the cross-sectional nature of our study does not permit one to infer causality. Fourth, self-report nature of the data should be kept in mind when interpreting some aspects such as medical attention and so on.

Conclusions

Results from this comparative study of suicidal thoughts and attempts in young adults from 12 Muslim countries may have several implications for science, prevention and policy. First, the percentages of students reporting nonfatal suicidal behavior are high despite prohibiting religio-cultural attitudes to suicide and show considerable cross-national variation. Therefore, the national health policy makers should consider developing national strategies for tackling this serious health problem. Second, having legal sanctions against suicide does not lessen the frequencies of suicidal behavior. Making suicide illegal may indeed prevent suicidal persons from seeking help when they are in an urgent need. Further, making it illegal may present suicide as a means for rebellion for youth. Third, asking whether someone has attempted suicide may be an ambiguous question and therefore should be followed by an inquiry about what happens after an attempt. Our results showed that 67% of those reporting suicide attempt said that nothing happened afterwards because it was not serious. Therefore, future scientific investigations should bear this in mind. Fourth, given the variations in the frequencies of self-reported suicidal thoughts and attempts between countries with similar cultural background, future studies may benefit from an examination of cultural meaning of suicidal behavior. Finally, further scientific investigations on suicidal behavior in the Muslim majority world are warranted.

Compliance with Ethical Standards

Conflict of Interest There is no conflict of interest for this paper.

Ethical Approval Ethical or IRB approval was obtained in all study sites (countries). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Since participation in the study was voluntary and full anonymity was guaranteed informed consent was not obtained from all individual participants. This was written in the first page of the survey.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. World Health Organisation. Preventing suicide: A global imperative. Geneva: WHO; 2014.
2. Goldman-Mellor SJ, Caspi A, Harrington H, Hogan S, Nada-Raja S, Poulton R, et al. Suicide attempt in young people: a signal for long-term health care and social needs. *JAMA Psychiatr*. 2014;71:119–27.
3. Shepard DS, Gurewich D, Lwin AK, Reed GA, Silverman MM. Suicide and suicidal attempts in the United States: costs and policy implications. *Suicide Life Threat Behav*. 2016;46:352–62.
4. Han B, Compton WM, Eisenberg D, Milazzo-Sayre L, McKeon R, Hughes A. Prevalence and mental health treatment of suicidal ideation and behavior among college students aged 18–25 years and their non-college-attending peers in the United States. *J Clin Psychiatry*. 2016;77:815–24.
5. Borges G, Benjet C, Orozco R, Medina-Mora ME, Menendez D. Alcohol, cannabis and other drugs and subsequent suicide ideation and attempt among young Mexicans. *J Psychiatr Res*. 2017;91:74–82.
6. Testa RJ, Michaels MS, Bliss W, Rogers ML, Balsam KF, Joiner T. Suicidal ideation in transgender people: gender minority stress and interpersonal theory factors. *J Abnorm Psychol*. 2017;126:125–36.
7. Miranda-Mendizábal A, Castellví P, Parés-Badell O, Almenara J, Alonso I, Blasco MJ, et al. Sexual orientation and suicidal behaviour in adolescents and young adults: systematic review and meta-analysis. *Br J Psychiatry*. 2017;211:77–87.
8. Björkenstam E, Kosidou K, Björkenstam C. Childhood household dysfunction and risk of self-harm: a cohort study of 107 518 young adults in Stockholm County. *Int J Epidemiol*. 2016;45:501–11.
9. Björkenstam E, Dalman C, Vinnerljung B, Weitoft GR, Walder DJ, Burström B. Childhood household dysfunction, school performance and psychiatric care utilisation in young adults: a register study of 96 399 individuals in Stockholm County. *J Epidemiol Community Health*. 2016;70:473–80.
10. Rostila M, Berg L, Arat A, Vinnerljung B, Hjern A. Parental death in childhood and self-inflicted injuries in young adults—a national cohort study from Sweden. *Eur Child Adolesc Psychiatry*. 2016;25:1103–11.
11. Rosiek A, Rosiek-Kryszewska A, Leksowski Ł, Leksowski K. Chronic stress and suicidal thinking among medical students. *Int J Environ Res Public Health*. 2016;13:212.
12. Buron P, Jimenez-Trevino L, Saiz PA, Garcia-Portilla MP, Corcoran P, Carli V, et al. Reasons for attempted suicide in Europe: prevalence, associated factors, and risk of repetition. *Arch Suicide Res*. 2016;20:45–58.
13. Kleiman EM, Liu RT. Social support as a protective factor in suicide: findings from two nationally representative samples. *J Affect Disord*. 2013;150:540–5.
14. Mackin DM, Perlman G, Davila J, Kotov R, Klein DN. Social support buffers the effect of interpersonal life stress on suicidal ideation and self-injury during adolescence. *Psychol Med*. 2017;47:1149–61.
15. Koch HJ. Suicides and suicide ideation in the Bible: an empirical survey. *Acta Psychiatr Scand*. 2005;112:167–72.
16. Nelson G, Hanna R, Hourii A, Klimes-Dougan B. Protective functions of religious traditions for suicide risk. *Suicidol Online*. 2012;3:59–71.
17. Eskin M, Voracek M, Stieger S, Altinyazar V. A cross-cultural investigation of suicidal behavior and attitudes in Austrian and Turkish medical students. *Soc Psychiatry Psychiatr Epidemiol*. 2011;46:813–23.
18. Eskin M. Gender and cultural differences in the 12-month prevalence of suicidal thoughts and attempts in Swedish and Turkish adolescents. *J Gend Cult Heal*. 1999;4:187–200.
19. Eskin M, Palova E, Krokavcova M. Suicidal behavior and attitudes in slovak and Turkish high school students: a cross-cultural investigation. *Arch Suicide Res*. 2014;18:58–73.
20. Eskin M, Sun J-M, Abuidhail J, Yoshimasu K, Kujan O, Janghorbani M, et al. Suicidal behavior and psychological distress in university students: a 12-nation study. *Arch Suicide Res*. 2016;20:369–88.
21. Lester D. Suicide and islam. *Arch Suicide Res*. 2006;10:77–97.
22. Pritchard C, Amanullah S. An analysis of suicide and undetermined deaths in 17 predominantly Islamic countries contrasted with the UK. *Psychol Med*. 2007;37:421–30.
23. Boehnke K, Lietz P, Schreier M, Wilhelm A. Sampling: the selection of cases for culturally comparative psychological research. Sampling: the selection of cases for culturally comparative psychological research. In: Matsumoto D, van de Vijver FJR, editors. *Culture and psychology. Cross-cultural research methods in psychology*. New York: Cambridge University Press; 2011. p. 101–29.
24. WHO. Suicide rates, age-standardized data by country. <http://apps.who.int/gho/data/node.main.MHSUICIDEASDR?lang=en>. 2015. Accessed 23 Feb 2017.
25. Behling O, Law KS. *Translating questionnaires and other research instruments: problems and solutions*. Thousand Oaks: Sage; 2000.
26. Boergers J, Spirito A, Donaldson D. Reasons for adolescent suicide attempts: associations with psychological functioning. *J Am Acad Child Adolesc Psychiatry*. 1998;37:1287–93.
27. Amiri L, Voracek M, Yousef S, Galadari A, Yammahi S, Sadeghi MR, et al. Suicidal behavior and attitudes among medical students in the United Arab Emirates. *Crisis*. 2013;34:116–23.

28. Eskin M. Suicidal behavior as related to social support and assertiveness among Swedish and Turkish high school students: a cross-cultural investigation. *J Clin Psychol.* 1995;51:158–72.
29. Chesney E, Goodwin GM, Fazel S. Risks of all-cause and suicide mortality in mental disorders: a meta-review. *World Psychiatry.* 2014;13:153–60.
30. Ferrari AJ, Norman RE, Freedman G, Baxter AJ, Pirkis JE, Harris MG, et al. The burden attributable to mental and substance use disorders as risk factors for suicide: findings from the global burden of disease study 2010. *PLoS One.* 2014;9:e91936.
31. Charara R, Forouzanfar M, Naghavi M, Moradi-Lakeh M, Afshin A, Vos T, et al. The burden of mental disorders in the eastern mediterranean region, 1990-2013. *PLoS One.* 2017;12:e0169575.
32. Garlow SJ, Rosenberg J, Moore JD, Haas AP, Koestner B, Hendin H, et al. Depression, desperation, and suicidal ideation in college students: results from the American Foundation for Suicide Prevention College screening project at Emory University. *Depress Anxiety.* 2008;25:482–8.
33. Wilcox HC, Arria AM, Caldeira KM, Vincent KB, Pinchevsky GM, O'Grady KE. Prevalence and predictors of persistent suicide ideation, plans, and attempts during college. *J Affect Disord.* 2010;127:287–94.
34. Sun L, Zhou C, Xu L, Li S, Kong F, Chu J. Suicidal ideation, plans and attempts among medical college students in China: the effect of their parental characteristics. *Psychiatry Res.* 2017;247:139–43.
35. Esen B, Gumuscu S. Rising competitive authoritarianism in Turkey. *Third World Q.* 2016;37:1581–606.
36. Hoffman M, Jamal A. The youth and the Arab spring: cohort differences and similarities. *Middle East Law Gov.* 2012;4:168–88.
37. Ciftci A, Jones N, Corrigan PW. Mental health stigma in the Muslim community. *J Muslim Ment Health.* 2013;7:17–32.
38. Pompili M, Goracci A, Giordano G, Erbuto D, Girardi P, Klonsky ED, et al. Relationship of non-suicidal self-injury and suicide attempt: a psychopathological perspective. *J Psychopathol.* 2015;21:348–53.
39. Ajdacic-Gross V, Weiss MG, Ring M, Hepp U, Bopp M, Gutzwiller F, et al. Methods of suicide: international suicide patterns derived from the WHO mortality database. *Bull World Health Organ.* 2008;86:726–32.
40. Stack S, Wasserman I. Race and method of suicide: culture and opportunity. *Arch Suicide Res.* 2005;9:57–68.
41. Nock MK. Self-injury. *Annu Rev Clin Psychol.* 2010;6:339–63.
42. Ward-Ciesielski EF, Schumacher JA, Bagge CL. Relations between nonsuicidal self-injury and suicide attempt characteristics in a sample of recent suicide attempters. *Crisis.* 2016;37:310–3.
43. Schrijvers DL, Bollen J, Sabbe BGC. The gender paradox in suicidal behavior and its impact on the suicidal process. *J Affect Disord.* 2012;138:19–26.
44. Värnik P. Suicide in the world. *Int J Environ Res Public Health.* 2012;9:760–71.
45. Denning DG, Conwell Y, King D, Cox C. Method choice, intent, and gender in completed suicide. *Suicide Life Threat Behav.* 2000;30:282–8.
46. Kleiman EM, Adams LM, Kashdan TB, Riskind JH. Grateful individuals are not suicidal: buffering risks associated with hopelessness and depressive symptoms. *Pers Individ Differ.* 2013;55:595–9.
47. Miller AB, Esposito-Smythers C, Leichtweis RN. Role of social support in adolescent suicidal ideation and suicide attempts. *J Adolesc Health.* 2015;56:286–92.

Mehmet Eskin, PhD, is a professor of clinical psychology at the department of Psychology, College of Social Sciences and Humanities, Koc University, Istanbul, Turkey. Dr. Eskin earned his bachelor's degree in psychology from the Department of Psychology, Middle East Technical University, in Ankara. He did his graduate training in clinical psychology in the Departments of Psychology in both Middle East Technical University and University of Oslo, Norway. He received his Ph.D. in psychology from the Department of Psychology, Stockholm University, Sweden. Dr. Eskin authored six books and nearly 100 scholarly articles on suicide, problem solving therapy and sexual orientation. He is an elected member of The Science Academy of Turkey.

Fadia AlBuhairan, MD, is a clinician/researcher/educator specialized in Pediatrics and Adolescent Medicine. She is a pioneer of Adolescent Medicine in the Arab world and has expertise in the complexity and management of adolescent and youth health issues. She established the first specialty adolescent health service and adolescent health research program in the region. She is also the first to incorporate and teach adolescent health into a medical school curriculum in Saudi Arabia. Currently she is Associate Professor in College of Medicine, at King Saud bin Abdulaziz University for Health Sciences, and Consultant Pediatrics and Adolescent Medicine in the Department of Pediatrics, King Abdulaziz Medical City, Ministry of the National Guard, Riyadh, Saudi Arabia. Her research interests are focused mainly in child and adolescent health.

Mohsen Rezaeian, PhD, got his in Epidemiology from Manchester University, England in 2002. Since then he works within the Epidemiology Department of Rafsanjan Medical School in Iran. His major interest lies in the epidemiology of suicide and its association with socio-economic status, cultural background, and religion. He is one of the founders of Iranian Scientific Society for Suicide Prevention (IRSSP) and the editor-in-chief of the newly developed Journal of Suicide Prevention (JSP). He is also appointed as the co-chair of Development of Effective National Suicide Prevention Strategy and Practice Special Interest Group of the International Association for Suicide Prevention (IASP). Epidemiology of natural disasters, wars and armed conflicts are among other his major interests. He publishes more than 300 articles in peer reviewed English and Persian journals.

Ahmed M. Abdel-Khalek, PhD, is a professor of Psychology in faculty of Arts, Alexandria University, Egypt, specialized in personality psychology and psychopathology. His research interests include the factorial study of personality dimensions, personality lexicon, cross-cultural comparisons in personality and intelligence, obsession-compulsion, anxiety, death anxiety, childhood depression, optimism and pessimism, religiosity, happiness, among others. He coined the terms and scales of death obsession and love of life.

Hacer Harlak, PhD, is a professor of social psychology and head of the Psychology Department at Adnan Menderes University in Aydin-Turkey. Dr. Harlak teaches social psychology and joint issues of social and clinical psychology in both undergraduate and graduate levels. Her published works focus on health behavior and attitudes, communication skills training, and intergroup relations.

Mayssah El-Nayal, PhD, is currently the Dean of the Faculty of Human Sciences, Head of the Psychology Department and Head of the Counselling Unit at Beirut Arab University. Between 2000 and 2007, Professor El-Nayal was the Head of the Psychology Department at Alexandria University, and from 1995 to 1997, she was the Head of the Mental Health Department at Qatar University. She obtained her BA in Psychology and MA in Social Psychology from Alexandria University. After that, she left to Goldsmith College at London University for 2 years to collect data for her PhD in Personality and Clinical Psychology which she later obtained from Alexandria University. Professor El-Nayal has several publications in national and international periodicals. She has always shown a great interest in conducting research, focusing on personality, clinical psychology and women' studies. Recently, her research has taken a new turn in the realm of positive psychology and the enhancement of its components in marginalized groups.

Nargis Asad, PhD, is an Associate Professor and Clinical Psychologist in the department of Psychiatry, Aga Khan University Karachi, Pakistan. She is also the chair of Working Group for Women (WGW) at AKU, the group works on gender issues through educational and research endeavors. Dr. Asad is actively involved in clinical and research work surrounding women/ adolescent mental health, domestic violence and suicide. Her research and clinical work is primarily multidisciplinary with colleagues at AKU and international partners.

Aqeel Khan, PhD, is faculty in the School of Education, Universiti Teknologi Malaysia (UTM) JOHOR, Malaysia. His current research interest includes Social Psychology, Positive Psychology and Educational Psychology. He has published 45 articles in Web of science Impact factor and Scopus indexed academic journals, contributed 17 chapters in research books and edited 6 books. At present he is executing research projects, supervising PhD and Master students and providing training for publications at national and international level. He has received many research excellence awards for his research and publications. Dr. Khan also teaching courses of research method and Counselling at post graduate level. He has presented many key notes addresses at various International Conferences at Singapore, Malaysia and Indonesia. He has travelled number of countries to pursue his academic interests like Indonesia, Thailand, India, South Africa, Lebanon and Singapore. He is also doing partnership in International projects in collaboration with Turkey, Indonesia, UK and Netherland.

Anwar Mechri, MD, is a professor of psychiatry in the Department of Psychiatry at the University Hospital of Monastir (Tunisia). Dr. Anwar Mechri is the Local primary researcher of the suicidal behavior and attitudes multinational project. His published works focus on schizophrenia and bipolar disorders and suicidal behaviors.

Isa Multazam Noor, MD, is a child psychiatrist at Dr. Soeharto Heerdjan Mental Hospital Jakara Indonesia. Dr. Noor received his medical education from Yarsi University Jakarta, psychiatric training from Gadjah Mada University Jogjakarta and child psychiatry residency training from University of Indonesia Jakarta. Child and adolescent mental health issues, traumatized children at abuse and natural disaster, psychosexual on child

milestones, biology psychiatry and psychopharmacology, family psychoeducation, religious/spirituality and psychiatry, suicide and depression prevention, sensory integration therapy, play therapy (especially on art), and community mental health on child and adolescents.

Motasem Hamdan, PhD, is the dean of Public Health, Al-Quds University. He got his bachelor and master's degrees in health management from Hacettepe University in Turkey and his PhD in Medical and Social Science (Health Policy) from the University of Leuven (KU Leuven) in Belgium. Dr. Hamdan is an expert in health system development. He served as senior health policy advisor to the Palestinian Ministry of Health and consultant for many national and international organizations. Dr. Hamdan is an author of more than 25 peer reviewed articles, member of editorial board of three internationally refereed journals and serves on a variety of local advisory boards. His research interests are in the areas of patient safety and quality of care, and health human resources development.

Ulker Isayeva, PhD Candidate, graduated from psychology department at Khazar University in Baku, Azerbaijan. She received her master's degree in Cognitive Neuroscience from York University, UK. In 2016, she has started pursuing a PhD degree at Koc University in Industrial and Organizational Psychology, Istanbul, Turkey. Her research interests include power distance, transformational leadership, women in leadership, fear of failure, etc. She currently continues her research activities both at Khazar University and Koc University.

Yousef Khader, BDS, MSc, MSPH, MHPE, FFPH, ScD, is a professor of Epidemiology and Biostatistics at Jordan University of Science and Technology (JUST). He has made substantial contributions to the understanding of the epidemiology of many diseases and health related problems in Jordan and in the region. He published more than 300 papers in reputable journals including the Lancet and New England Journal of Medicine. He is the principal investigator of many ongoing funded studies.

Alaa Al Sayyari, MSc. graduated from the college of Applied Medical Science at King Saud University, Riyadh, Saudi Arabia, and completed her master's Degree of Health Care Management at California State University, Los Angeles, USA. She currently works in Population-Health research at King Abdullah International Medical Research Center, Riyadh, Saudi Arabia, with an interest in the field of substance abuse and mental health.

Albaraa Khader, is a sixth-year medical student at Jordan University of Science and Technology. He is involved in many research projects in Jordan and he is the research coordinator for the National Perinatal Mortality Study in Jordan (J-SANDS).

Bahareh Behzadi is a Master student of epidemiology within the Epidemiology Department of Rafsanjan Medical School in Iran. She assists Professor Mohsen Rezaeian on "suicidal behavior among students in Muslim countries" Project.

Cennet Şafak Öztürk, PhD, earned her BSc, MSc and Doctoral degrees from the psychology department of Ege University in Izmir, Turkey. Currently she is the assistant professor of clinical psychology at Adnan Menderes University Faculty of Arts and Sciences, Department of Psychology, Aydin, Turkey. Her research interests include sexuality, sexual Function and sexual dysfunction, personality, and psychotherapy.

Laifa Hendarmin, PhD, is the head of Medical Research Unit in Faculty of Medicine, Syarif Hidayatullah State Islamic University in Jakarta, Indonesia. Since 2014, she has joined Center for the Study of Islam and Society (PPIM), the most leading research center in the university. She graduated from Faculty of Dentistry, University of Indonesia in 2002, and gained her Ph.D. from Graduate School of Dental Science, Kyushu University, Japan, in 2008. Having a basic science background, she currently enriches her research interest with social approaches, including smoking and vaping behavior in many communities, such as students in Islamic boarding school and HIV-infected drug users. She also involves in many other topics of health and social studies, specially related with young age, including youth risk behavior, suicide ideas, Internet addiction, radical opinions and actions amongst Muslim students.

Murad Moosa Khan, MBBS, MR PSYCH, PhD is Professor, Dept. of Psychiatry, Aga Khan University, Karachi, Pakistan. He is the current President of the International Association for Suicide Prevention (IASP). His clinical & research interests include suicide & self-harm, mental health of women & elderly & medical ethics.

Salam Khatib, PhD, is the coordinator of master studies at nursing department at AlQuds University. Her research interests are focused in different areas: (a) honor killing, (b) postpartum depression and attachment, (c) family planning, (d) Type 1 diabetes among children (e) Congenital heart disease among children and stress and burn out among oncology nurses. An important theme in her work is gender-based violence and particularly honor killing. She has an interest in qualitative research such as grounded theory and content analysis.

Affiliations

Mehmet Eskin¹ • Fadia AlBuhairan^{2,3} • Mohsen Rezaeian⁴ • Ahmed M. Abdel-Khalek⁵ • Hacer Harlak⁶ • Mayssah El-Nayal⁷ • Nargis Asad⁸ • Aqeel Khan⁹ • Anwar Mechri¹⁰ • Isa Multazam Noor¹¹ • Motasem Hamdan¹² • Ulker Isayeva¹³ • Yousef Khader¹⁴ • Alaa Al Sayyari¹⁵ • Albaraa Khader¹⁴ • Bahareh Behzadi⁴ • Cennet Şafak Öztürk⁶ • Laifa Annisa Hendarmin¹⁶ • Murad Moosa Khan⁸ • Salam Khatib¹⁷

¹ Department of Psychology, College of Social Sciences and Humanities, Koc University, Rumelifeneriyolu 34450 Sariyer, Istanbul, Turkey

² Al Dara Hospital and Medical Center, Riyadh, Saudi Arabia

³ Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

⁴ Epidemiology and Biostatistics, Rafsanjan Medical School, Rafsanjan, Iran

⁵ Department of Psychology, Faculty of Arts, Alexandria University, Alexandria, Egypt

⁶ Department of Psychology, Faculty of Arts and Sciences, Adnan Menderes University, Aydin, Turkey

⁷ Department of Psychology, Faculty of Human Sciences, Beirut Arab University, Beirut, Lebanon

⁸ Department of Psychiatry, Medical College, Aga Khan University, Karachi, Pakistan

⁹ Faculty of Education, Universiti Teknologi Malaysia, Johor, Malaysia

¹⁰ Department of Psychiatry, University hospital of Monastir, Monastir, Tunisia

¹¹ Dr Soeharto Heerdjan Mental Hospital, Jakarta, Indonesia

¹² School of Public Health, Al-Quds University, Jerusalem, Palestine

¹³ Department of Psychology, Khazar University, Baku, Azerbaijan

¹⁴ Department of Community Medicine, Public Health and Family Medicine, Faculty of Medicine, Jordan University of Science & Technology, Irbid, Jordan

¹⁵ Population Health Research Section-Hospital-MNGHA, King Abdullah International Medical Research Center / King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

¹⁶ Faculty of Medicine, Syarif Hidayatullah State Islamic University, Jakarta, Indonesia

¹⁷ Faculty of Health Professions, Department of Nursing, Al-Quds University, Jerusalem, Palestine